

SPECIALTY GUIDELINE MANAGEMENT

Makena (hydroxyprogesterone caproate injection)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered covered benefits provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indication

Makena is indicated to reduce the risk of preterm birth in women with a singleton pregnancy who have a history of singleton spontaneous preterm birth. The effectiveness of Makena is based on improvement in the proportion of women who delivered < 37 weeks of gestation. There are no controlled trials demonstrating a direct clinical benefit, such as improvement in neonatal mortality and morbidity.

Limitations of use: While there are many risk factors for preterm birth, safety and efficacy of Makena has been demonstrated only in women with a prior spontaneous singleton preterm birth. **It is not intended for use in women with multiple gestations or other risk factors for preterm birth.**

All other indications are considered experimental/investigational and are not covered benefits.

II. EXCLUSIONS

Coverage will not be provided for members with any of the following exclusions:

- A. Current or history of thrombosis or thromboembolic disorders
- B. Known or suspected breast cancer, other hormone-sensitive cancer, or a history of these conditions
- C. Undiagnosed abnormal vaginal bleeding unrelated to pregnancy
- D. Cholestatic jaundice of pregnancy
- E. Liver tumors, benign or malignant, or active liver disease
- F. Uncontrolled hypertension

III. CRITERIA FOR INITIAL APPROVAL

Prevention Of Preterm Birth

Authorization of 21 weeks or through 36 weeks, 6 days of gestational age, whichever is less, may be granted for the prevention of preterm birth when all of the following criteria are met:

- A. The current pregnancy is a singleton pregnancy.
- B. The member has a history of singleton spontaneous preterm birth, defined as delivery at less than 37 weeks gestation following preterm labor, preterm rupture of membranes, and cervical insufficiency.
- C. Makena will be initiated between 16 weeks, 0 days and 24 weeks, 6 days of gestation.

Makena SGM P2017x

CVS Caremark is an independent company that provides pharmacy benefit management services to CareFirst BlueCross BlueShield and CareFirst BlueChoice, Inc. members.

CareFirst BlueCross BlueShield is the shared business name of CareFirst of Maryland, Inc. and Group Hospitalization and Medical Services, Inc. CareFirst of Maryland, Inc., Group Hospitalization and Medical Services, Inc., CareFirst BlueChoice, Inc., The Dental Network and First Care, Inc. are independent licensees of the Blue Cross and Blue Shield Association. In the District of Columbia and Maryland, CareFirst MedPlus is the business name of First Care, Inc. In Virginia, CareFirst MedPlus is the business name of First Care, Inc. of Maryland (used in VA by: First Care, Inc.). © Registered trademark of the Blue Cross and Blue Shield Association

IV. CONTINUATION OF THERAPY

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

V. REFERENCES

1. Makena [package insert]. Waltham, MA: Lumara Health Inc.; April 2016.
2. American College of Obstetricians and Gynecologists Committee on Practice Bulletins—Obstetrics. ACOG practice bulletin no. 130: prediction and prevention of preterm birth. *Obstet Gynecol.* 2012;120(4):964-973.