

SPECIALTY GUIDELINE MANAGEMENT

RASUVO (methotrexate injection)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

1. Rheumatoid Arthritis including Polyarticular Juvenile Idiopathic Arthritis
Rasuvo is indicated in the management of selected adults with severe, active rheumatoid arthritis (RA) or children with active polyarticular juvenile idiopathic arthritis, who have had an insufficient therapeutic response to, or are intolerant of, an adequate trial of first-line therapy including full dose non-steroidal anti-inflammatory agents (NSAIDs).
2. Psoriasis
Rasuvo is indicated in adults for the symptomatic control of severe, recalcitrant, disabling psoriasis that is not adequately responsive to other forms of therapy, but only when the diagnosis has been established, as by biopsy and/or after dermatologic consultation. It is important to ensure that a psoriasis “flare” is not due to an undiagnosed concomitant disease affecting immune responses.

All other indications are considered experimental/investigational and are not a covered benefit.

II. CRITERIA FOR INITIAL APPROVAL

Authorization of 24 months may be granted for treatment of RA, pJIA, or psoriasis when BOTH of the following criteria are met:

- A. Member has tried and had an inadequate response or intolerance to generic oral methotrexate.
- B. Member has inability to prepare and administer generic injectable methotrexate.

III. CONTINUATION OF THERAPY

Authorization of 24 months may be granted for all members (including new members) who meet ALL initial authorization criteria and achieve or maintain positive clinical response after at least 3 months of therapy with Rasuvo as evidenced by low disease activity or improvement in signs and symptoms of the condition.

IV. REFERENCES

1. Rasuvo [package insert]. Chicago, IL: Medac Pharma Inc.; November 2014.

2. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol.* 2016;68(1):1-26.
3. Smolen JS, Landewé R, Billsma J, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis.* 2017;0:1-18.
4. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring of therapeutic agents for the treatment of arthritis and systemic features. *Arthritis Care Res.* 2011;63(4):465-482.
5. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4: Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. *J Am Acad Dermatol.* 2009;61:451-485.