

SPECIALTY GUIDELINE MANAGEMENT

Supprelin LA (histrelin acetate)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indication

Supprelin LA is indicated for the treatment of children with central precocious puberty.

B. Compendial Use

Gender Dysphoria (also known as gender non-conforming or transgender persons)

NOTE: Some plans may opt-out of coverage for gender dysphoria.

All other indications are considered experimental/investigational and are not a covered benefit.

II. CRITERIA FOR INITIAL APPROVAL

A. Central Precocious Puberty (CPP)

1. Authorization up to age 12 may be granted for the treatment of CPP in a female member when ALL of the following criteria are met:
 - a. The diagnosis of CPP has been confirmed by a pubertal response to a gonadotropin releasing hormone (GnRH) agonist test or a pubertal level of a third generation luteinizing hormone (LH) assay
 - b. The diagnosis of CPP has been confirmed by assessment of bone age versus chronological age
 - c. The member was less than 8 years of age at the onset of secondary sexual characteristics
2. Authorization up to age 13 may be granted for the treatment of CPP in a male member when ALL of the following criteria are met:
 - a. The diagnosis of CPP has been confirmed by a pubertal response to a GnRH agonist test or a pubertal level of a third generation LH assay
 - b. The diagnosis of CPP has been confirmed by assessment of bone age versus chronological age
 - c. The member was less than 9 years of age at the onset of secondary sexual characteristics

B. Gender Dysphoria

1. Authorization of 12 months may be granted for pubertal suppression in preparation for gender reassignment in an adolescent member when ALL of the following criteria are met:
 - a. The member has a diagnosis of gender dysphoria
 - b. The member has reached Tanner stage 2 of puberty

Supprelin LA SGM P2017a

CVS Caremark is an independent company that provides pharmacy benefit management services to CareFirst BlueCross BlueShield and CareFirst BlueChoice, Inc. members.

CareFirst BlueCross BlueShield is the shared business name of CareFirst of Maryland, Inc. and Group Hospitalization and Medical Services, Inc. CareFirst of Maryland, Inc., Group Hospitalization and Medical Services, Inc., CareFirst BlueChoice, Inc., The Dental Network and First Care, Inc. are independent licensees of the Blue Cross and Blue Shield Association. In the District of Columbia and Maryland, CareFirst MedPlus is the business name of First Care, Inc. In Virginia, CareFirst MedPlus is the business name of First Care, Inc. of Maryland (used in VA by: First Care, Inc.). © Registered trademark of the Blue Cross and Blue Shield Association

2. Authorization of 12 months may be granted for gender reassignment in an adult member when ALL of the following criteria are met:
 - a. The member has a diagnosis of gender dysphoria
 - b. The member will receive Supprelin LA concomitantly with cross sex hormones

III. CONTINUATION OF THERAPY

A. CPP

1. Authorization up to age 12 may be granted for continuation of therapy for CPP in a female member if the member is currently less than 12 years of age.
2. Authorization up to age 13 may be granted for continuation of therapy for CPP in a male member if the member is currently less than 13 years of age.

B. Gender Dysphoria

All members (including new members) requesting authorization for continuation of therapy must meet ALL initial authorization criteria.

IV. REFERENCES

1. Supprelin LA [package insert]. Malvern, PA: Endo Pharmaceuticals Solutions Inc.; June 2013.
2. Kletter GB, Klein KO, Wong YY. A pediatrician's guide to central precocious puberty. *Clin Pediatr*. 2015;54:414-424.
3. Carel J, Eugster EA, Rogol A, et al. Consensus statement on the use of gonadotropin-releasing hormone analogs in children. *Pediatrics*. 2009;123:e752-e762.
4. Houk CP, Kunselman AR, Lee PA. Adequacy of a single unstimulated luteinizing hormone level to diagnose central precocious puberty in girls. *Pediatrics*. 2009;123:e1059-e1063.
5. Kaplowitz P, Bloch C, the Section on Endocrinology. Evaluation and referral of children with signs of early puberty. *Pediatrics*. 2016;137:e20153732.
6. Hembree WC, Cohen-Kettenis P, Delemarre-van de Waal HA, et al. Endocrine treatment of transsexual persons: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2009;94:3152-3154.
7. Gender Identity Research and Education Society. Guidance for GPs and other clinicians on the treatment of gender variant people. UK Department of Health. Published March 10, 2008.
8. Standards of care for the health of transsexual, transgender, and gender-nonconforming people, 7th version. ©2012 World Professional Association for Transgender Health. Available at <http://www.wpath.org>.