

## SPECIALTY GUIDELINE MANAGEMENT

### TRACLEER (bosentan)

#### POLICY

#### I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

##### FDA-Approved Indication

##### **Pulmonary Arterial Hypertension (PAH)**

Tracleer is indicated for the treatment of PAH (WHO Group 1):

- A. In adults to improve exercise ability and to decrease clinical worsening.
- B. In pediatric patients aged 3 years and older with idiopathic or congenital PAH to improve pulmonary vascular resistance (PVR), which is expected to result in an improvement in exercise ability.

All other indications are considered experimental/investigational and are not a covered benefit.

#### II. CRITERIA FOR INITIAL APPROVAL

Authorization of 12 months may be granted for treatment of PAH when ALL of the following criteria are met:

- A. Member has PAH defined as WHO Group 1 class of pulmonary hypertension (refer to Appendix).
- B. PAH was confirmed by either criterion (1) or criterion (2) below:
  1. Pretreatment right heart catheterization with all of the following results:
    - mPAP  $\geq$  25 mmHg
    - PCWP  $\leq$  15 mmHg
    - PVR  $>$  3 Wood units
  2. For infants less than one year of age with any of the following conditions, PAH was confirmed by Doppler echocardiogram if right heart catheterization cannot be performed:
    - Post cardiac surgery
    - Chronic heart disease
    - Chronic lung disease associated with prematurity
    - Congenital diaphragmatic hernia

#### III. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for members with PAH who are currently receiving Tracleer therapy through a paid pharmacy or medical benefit.

#### IV. APPENDIX

##### **WHO Classification of Pulmonary Hypertension**

##### WHO Group 1. Pulmonary Arterial Hypertension (PAH)

##### 1.1 Idiopathic (IPAH)

Tracleer SGM P2017

CVS Caremark is an independent company that provides pharmacy benefit management services to CareFirst BlueCross BlueShield and CareFirst BlueChoice, Inc. members.

CareFirst BlueCross BlueShield is the shared business name of CareFirst of Maryland, Inc. and Group Hospitalization and Medical Services, Inc. CareFirst BlueCross BlueShield and CareFirst BlueChoice, Inc. are both independent licensees of the Blue Cross and Blue Shield Association. The Blue Cross and Blue Shield Names and Symbols are registered trademarks of the Blue Cross and Blue Shield Association. © Registered trademark of CareFirst of Maryland, Inc.

- 1.2 Heritable PAH
  - 1.2.1 Germline mutations in the bone morphogenetic protein receptor type 2 (BMP2)
  - 1.2.2 Activin receptor-like kinase type 1 (ALK1), endoglin (with or without hereditary hemorrhagic telangiectasia), Smad 9, caveolin-1 (CAV1), potassium channel super family K member-3 (KCNK3)
  - 1.2.3 Unknown
- 1.3 Drug- and toxin-induced
- 1.4. Associated with:
  - 1.4.1 Connective tissue diseases
  - 1.4.2 HIV infection
  - 1.4.3 Portal hypertension
  - 1.4.4 Congenital heart diseases
  - 1.4.5 Schistosomiasis
- 1'. Pulmonary veno-occlusive disease (PVOD) and/or pulmonary capillary hemangiomatosis (PCH)
- 1". Persistent pulmonary hypertension of the newborn (PPHN)

WHO Group 2. Pulmonary Hypertension Owing to Left Heart Disease

- 2.1 Systolic dysfunction
- 2.2 Diastolic dysfunction
- 2.3 Valvular disease
- 2.4 Congenital/acquired left heart inflow/outflow tract obstruction and congenital cardiomyopathies

WHO Group 3. Pulmonary Hypertension Owing to Lung Disease and/or Hypoxia

- 3.1 Chronic obstructive pulmonary disease
- 3.2 Interstitial lung disease
- 3.3 Other pulmonary diseases with mixed restrictive and obstructive pattern
- 3.4 Sleep-disordered breathing
- 3.5 Alveolar hypoventilation disorders
- 3.6 Chronic exposure to high altitude
- 3.7 Developmental abnormalities

WHO Group 4. Chronic Thromboembolic Pulmonary Hypertension (CTEPH)

WHO Group 5. Pulmonary Hypertension with Unclear Multifactorial Mechanisms

- 5.1 Hematologic disorders: Chronic hemolytic anemia, myeloproliferative disorders, splenectomy
- 5.2 Systemic disorders: sarcoidosis, pulmonary Langerhans cell histiocytosis: lymphangioleiomyomatosis, neurofibromatosis, vasculitis
- 5.3 Metabolic disorders: glycogen storage disease, Gaucher disease, thyroid disorders
- 5.4 Others: tumoral obstruction, fibrosing mediastinitis, chronic renal failure on dialysis, segmental PH

## V. REFERENCES

1. Tracleer [package insert]. South San Francisco, CA: Actelion Pharmaceuticals US, Inc.; September 2017.
2. Chin KM, Rubin LJ. Pulmonary arterial hypertension. *J Am Coll Cardiol.* 2008;51(16):1527-1538.
3. McLaughlin VV, Archer SL, Badesch DB, et al. ACCF/AHA 2009 expert consensus document on pulmonary hypertension a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association developed in collaboration with the American College of Chest Physicians; American Thoracic Society, Inc.; and the Pulmonary Hypertension Association. *J Am Coll Cardiol.* 2009;53(17):1573-1619.
4. Badesch DB, Champion HC, Gomez-Sanchez MA, et al. Diagnosis and assessment of pulmonary arterial hypertension. *J Am Coll Cardiol.* 2009;54:S55-S66.
5. Simonneau G, Robbins IM, Beghetti M, et al. Updated clinical classification of pulmonary hypertension. *J Am Coll Cardiol.* 2013;62:D34-S41.
6. Rubin LJ; American College of Chest Physicians. Diagnosis and management of pulmonary arterial hypertension: ACCP evidence-based clinical practice guidelines. *Chest.* 2004;126(1 Suppl):7S-10S.
7. Barst RJ, Gibbs SR, Ghofrani HA, et al. Updated evidence-based treatment algorithm in pulmonary arterial hypertension. *J Am Coll Cardiol.* 2009;54:S78-S84.
8. Taichman DB, Ornelas J, Chung L, et al. Pharmacologic therapy for pulmonary arterial hypertension in adults. CHEST guideline and expert panel report. *Chest.* 2014;46(2):449-475.
9. Abman, SH, Hansmann G, Archer SL, et al. Pediatric pulmonary hypertension: guidelines from the American Heart Association and American Thoracic Society. *Circulation.* 2015;132(21):2037-99.