



SPECIALTY GUIDELINE MANAGEMENT

SYMDEKO (tezacaftor/ivacaftor)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indication

Symdeko is a combination of tezacaftor and ivacaftor, indicated for the treatment of patients with cystic fibrosis (CF) aged 12 years and older who are homozygous for the *F508del* mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor based on *in vitro* data and/or clinical evidence.

If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.

All other indications are considered experimental/investigational and are not a covered benefit.

II. REQUIRED DOCUMENTATION

The following information is necessary to initiate the prior authorization review: genetic testing report confirming the presence of the appropriate *CFTR* gene mutation.

III. CRITERIA FOR INITIAL APPROVAL

Cystic Fibrosis

Indefinite authorization may be granted for treatment of cystic fibrosis when all of the following criteria are met:

- Genetic testing was conducted to detect a mutation in the CFTR gene.
- b. The member has one of the following mutations in the *CFTR* gene: A455E, A1067T, D110E, D110H, D579G, D1152H, D1270N, E56K, E193K, E831X, F1052V, F1074L, K1060T, L206W, P67L, R74W, R117C, R347H, R352Q, R1070W, S945L, S977F, 711+3A→G, 2789+5G→A, 3272-26A→G, 3849+10kbC→T, or the member is homozygous for the F508del mutation.
- c. The member is at least 12 years of age.
- d. Symdeko will not be used in combination with Kalydeco or Orkambi.

IV. CONTINUATION OF THERAPY

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

Symdeko SGM P2018

V. REFERENCES

- Symdeko [package insert]. Boston, MA: Vertex Pharmaceuticals Inc.; February 2018.
 Rowe SM, Daines C, Ringshausen FC, Kerem E, Wilson J, Tullis E, Nair N, Simard C, Han L, Ingenito EP, McKee C, Lekstrom-Himes J, Davies JC. Tezacattor-Ivacaftor in Residual Funtion Heterzygotes with Cystic Fibrosis. N Engl J Med. 2017; 377:2024-2035
- Taylor-Cousar JL, Munck A, McKone EF, et al. Tezacaftor–ivacaftor in patients with cystic fibrosis 3. homozygous for Phe508del N Engl J Med 2017; 377:2013-2023